

Amended Claims

1. (three times amended) A method for inhibiting metastasis of lymphoma in a mammal which comprises administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma.

22. (twice amended) A method for treating lymphoma in a mammal which comprises subcutaneously administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma.

24. (twice amended) A method for treating lymphoma in a mammal which comprises intracavitarily administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma.

25. (twice amended) A method for treating [metastatic] lymphoma in a mammal which comprises administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma, wherein the method further comprises subjecting the mammal to at least one other treatment modality selected from the group consisting of immunotherapy, radiation therapy and surgery, and wherein the other treatment modality is administered prior to, during or after the administration of IVIG or fragments thereof.

26. (twice amended) A method for treating [metastatic] Hodgkin's lymphoma in a mammal which comprises administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂,

Fab', Fab and Fc, wherein the mammal has metastatic lymphoma, wherein the method further comprises subjecting the mammal to at least one other treatment modality.

28. (twice amended) A method for treating [metastatic] lymphoma in a mammal which comprises administering to the mammal a preparation of IVIG or fragments thereof selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma, wherein the method further comprises surgically removing the lymphoma prior to, during or after the administration of the preparation of IVIG or fragments thereof.

29. (twice amended) A method for treating [metastatic] lymphoma in a mammal which comprises administering to the mammal a preparation of fragments of IVIG selected from the group consisting of F(ab')₂, Fab', Fab and Fc, wherein the mammal has metastatic lymphoma.